

EO for Understanding and Monitoring Changing Conditions in Tropical Peatlands



Michel Bechtold



Frieke Van Coillie



Patrick Willems







Raphael Tshimanga







+ 2 PhD (Belgium) + PhD / technical staff (DRC and Singapore)





Peat ≈ 90% porosity

• Threats: Anthropogenic drainage and climate change

Peatland distribution



peatland distribution

peat in soil mosaic

Boundaries: United Nations Geospatial, 2021. The boundaries and names shown, and the designations used on this map do not imply official endorsement or acceptance by the United Nations. Peatland distribution: Global Peatland Database, 2022. Elevation: Jarvis et al. 2008. SRTM for the globe version 4.

EO4PEAT domains



~2% of annual anthropogenic CO_2 emissions




initial phase of disturbance








state-of-the-art




- Water table depth is main control of GHG balance in peatlands
- Strong connection between vegetation and hydrology



Peatland-specific data assimilation product
EO-based mapping of peatland land use land cover change (LULCC)




- Enhancing the accuracy of peatland-specific monitoring
- Improving the process understanding in peatlands facing different types of human and climate disturbance










*LULCC: Land use land cover change



WP1: inundation monitoring

Goal: Monitoring inundation dynamics using L-band active microwave data




- GNSS-R and SAR
- Statistical vs. physics-based retrievals
 → use of full-wave radar modeling
- Validation:
 - e.g., Brunei flux tower (International partner: Cobb)






Goal: Near real-time peatland-specific LULCC monitoring



2022, Sc. Data



Water Seasonal water Pristine peat swamp forest Degraded peat swamp forest Tall shrub/secondary forest Ferns/low shrub Small-holder area Industrial plantations Built-up area Clearance Mangrove • Fully convolutional neural networks


WP2: LULCC monitoring

- Use of 3D convolutional layers to capture both spatial and temporal features
- \rightarrow Time series of class probabilities



WP3: river-peatland interaction


Goal: Monitoring and modeling of river-peatland interaction






WP4: data assimilation

Goal: Gaining accuracy and insights from data assimilation with parameter updating





Data assimilation without (or with low order) of prior rescaling of observations
 → Reduce spatially variable model bias




WP5: human and climate disturbance


Goal: Attributing changing conditions to the cause of disturbance

• Analysis of abrupt changes or trends in hydrology and vegetation

 Scenario-based separation of climate-related trends and direct human induced trends










Peat Sarah Chadburn ♂ Angela Gallego-Sala ♂ Noah Smith ♂ Michel Bechtold ♂



expected impact

Innovative exploitation of EO data for peatlands

 New datasets and tools (In situ data, sub-canopy inundation, LULCC, modeling and DA)



- Insights into changes in hydrology and vegetation and their interplay
- Communication to academic and policy/broad public sector for optimized climate action





research team

KU LEUVEN



Peatland research Land surface modeling Data assimilation

Frieke Van Coillie



Object-Based Image Analysis / LULCC **Remote Sensing**

Raphael Tshimanga



Monitoring and modeling of Congo basin hydrology



Patrick Willems

River hydrology and hydraulics Regional hydrological modeling

Sébastien Lambot



UCLouvain

Radar electromagnetic modeling Inverse modeling



Alex Cobb

Peatland research Monitoring and modeling of Southeast Asian peatlands

Thanks for your attention!



Extra slides

04PEAT NASA E Quantifying ground/surface water influence Bangui river stage 8 total water storage (TWS) 1 0.8 (mm) comedian? increments from SMOS DA. С 40 0.6 Pearson Increments 0.4 0 0.2 40 0.0 -0.2 80 Climatology Stage height (m) -0.410 ი -0.6ω In-situ -0.8 ŝ DA diagnostics (SMOS) suggest influence of river stage height anomalies on peatland water tables Logging Concession Oil Palm Concession

• possible drainage

100 200 300 km

Peatland

Other Landcover Class



research team



Michel Bechtold | Gabrielle De Lannoy



Frieke Van Coillie









Sébastien Lambot



Raphael Tshimanga





Thanks for your attention!